



US006706753B2

(12) **United States Patent**
Artis et al.

(10) **Patent No.:** **US 6,706,753 B2**
(15) **Date of Patent:** **Mar. 16, 2004**

(54) **INTEGRIN RECEPTOR INHIBITORS**

(75) **Inventors:** **Dean R. Artis, Kensington, CA (US); David Y. Jackson, San Bruno, CA (US); Thomas E. Rawson, Mountain View, CA (US); Mark E. Reynolds, Millbrae, CA (US); Daniel P. Sutherland, South San Francisco, CA (US); Mark S. Stanley, Pacifica, CA (US)**

(73) **Assignee:** **Genentech, Inc., South San Francisco, CA (US)**

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl No** 10/313,147

(22) **Filed:** **Dec. 6, 2002**

(65) **Prior Publication Data**

US 2003/0100599 A1 May 29, 2003

Related U.S. Application Data

(63) Continuation of application No. 09/932,695, filed on Aug. 16, 2001, now abandoned

(60) Provisional application No. 60/226,626, filed on Aug. 18, 2000.

(51) **Int. Cl.** **A61K 31/4025; C07D 207/04**

(52) **U.S. Cl.** **514/422, 548/517; 548/537; 548/538, 514/423**

(58) **Field of Search** **548/517, 537, 548/538, 514/422, 423**

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,079,260 A 1/1992 Weitzberg et al.
5,223,517 A 6/1993 Muller et al. 514/339
5,272,161 A 12/1993 Niewolner et al.
5,334,606 A 8/1994 MacLeod
5,519,043 A 5/1996 Perumaram
5,773,465 A 6/1998 Gandolfi et al.
5,773,446 A 6/1998 Chandrakumar et al.
5,891,899 A 4/1999 Clark et al.
5,908,863 A 6/1999 Allegretti et al.
5,994,351 A 11/1999 Robinson et al. 514/237.5

FOREIGN PATENT DOCUMENTS

EP 327877 8/1989
JP 480540n1 7/1973
WO WO 97/01540 1/1997
WO WO 99/10312 3/1999

OTHER PUBLICATIONS

Andrew, H. E., et al., "Nitration of 3-Methylfluoranthene" *J. Chem. Soc., Perkins Trans. 1* vol. 6:755-759 (1972)
Burcher et al., "Lymphocyte Homing and Homeostasis" *Science* 272:60-66 (1996).

Cardarelli et al., "Cyclic RGD Peptide Inhibits $\alpha 4 \beta 1$ Interaction with Connecting Segment 1 and Vascular Cell Adhesion Molecule" *Journal of Biological Chemistry* 269:18668-18673 (1994).

Chulnyan et al., " $\alpha 4$ -Integrin-dependent emigration of monocytes" *Springer Semin. Immunopathology* 16:391-404 (1995).

Ehlers et al., "VCAM-1 on activated endothelium interacts with the leukocyte integrin VLA-4 at a site distinct from the VLA-4/fibronectin binding site" *Cell* 60:577-584 (1990).

Fotouhi et al., "Cyclic Thioether Peptide Mimetics as VCAM-VLA-4 Antagonists" *Bioorganic & Medicine Chemistry Letters* 10:1167-1169 (2000).

Hamilton, et al., "Fluorenylalkanoic and Benzoic Acids as Novel Inhibitors of Cell Adhesion Processes in Leukocytes" *J. Med. Chem.* 38:1650-1656 (1995).

Hemler, "VLA proteins in the integrin family: structures, functions, and their role on leukocytes" *Annu. Rev. Immunol.* 8:365-400 (1990).

Hynes, "Integrins: versatility, modulation, and signaling in cell adhesion" *Cell* 69(1):11-25 (1992).

Jackson et al., "Potent $\alpha 4 \beta 1$ Peptide Antagonists as Anti-Inflammatory Agents" *Journal of Medicinal Chemistry* 40(21):3359-3368 (Oct. 10, 1997).

Juliano et al., "Adhesion molecules in cancer: the role of integrins" *Current Opinion Cell Biology* 5:812-818 (1993).

Laberge, S. et al., "Role of VLA-4 and LFA-1 in Allergen-Induced Airway Hyperresponsiveness and Lung Inflammation in the Rat" *Am. J. Respir. Crit. Care Med.* 151:822-829 (1995).

Makarewicz et al., "Competitive binding of vascular cell adhesion molecule-1 and the HepII/III/CS domain of fibronectin to the integrin $\alpha 4 \beta 1$ " *Journal of Biological Chemistry* 269:4005-4011 (1994).

(List continued on next page.)


Primary Examiner—Joseph K. McKane

Assistant Examiner—Gulam M M Snameem

(74) **Attorney, Agent, or Firm**—David W Evans

(57) **ABSTRACT**

Provided are compounds of formula (I)



wherein A, Q, W, X, Y, Z, R₁ to R_n, m and n are as defined herein. Compounds of the invention bind to α_n integrin receptors and thereby inhibit binding of ligands for α_n integrins which is useful for prophylactic and/or therapeutic treatment of diseases and conditions associated with α_n integrins or their ligands

10 Claims, No Drawings

US 6,706,753 B2

Page 2

OTHER PUBLICATIONS

Nakajima et al., "Role of vascular cell adhesion molecule 1/very late activation antigen 4 and intercellular adhesion molecule 1/lymphocyte function-associated antigen 1 interactions in antigen-induced eosinophil and T cell recruitment into the tissue" *Journal of Experimental Medicine*.

Sbarat et al., "The adhesion cascade and anti-adhesion therapy: an overview" *Springer Semin. Immunopathology* 16 359-378 (1995)

Simmons et al., "Vascular cell adhesion molecule-1 expressed by bone marrow stromal cells mediates the binding of hematopoietic progenitor cells" *Blood* 80:388-395 (1992)

Database Crossfire Beilstein 'Online', Beilstein Institut zur Forderung der Chemischen Wissenschaften, Frankfurt am Main, DE; Database accession No. 2187180 XP002194815 Abstract, & Campbell, Crombie: Proc. Royal Soc. London, Series A, Mathematical & Physical Sciences, 65:376 (1960-1961) Royal Society of.

US 6,706,753 B2

1
INTEGRIN RECEPTOR INHIBITORS

RELATED APPLICATIONS

This application is a continuation application filed under 37 CFR 1.53(b)(1), claiming priority under 35 USC §120 to application Ser. No. 09/932,695, filed Aug. 16, 2001 now abandoned, and under 35 USC §119(e) to provisional application Serial No. 60/226,626, filed Aug. 18, 2000, the contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to novel compounds useful as therapeutic, prophylactic or diagnostic agents having binding affinity to integrin receptors, in particular to α_x integrins.


BACKGROUND OF THE INVENTION

The integrins are $\alpha\beta$ heterodimeric cell surface receptors involved in numerous cellular processes from cell adhesion to gene regulation (Hynes, Cell 1992, 69, 11-25; Hemler, Annu. Rev. Immunol. 1990, 8, 365-368). Several integrins have been implicated in disease processes and have generated widespread interest as potential targets for drug discovery (Sharar et al, Springer Semin. Immunopathology 1995, 16, 359-378). In the immune system, integrins are involved in leukocyte trafficking, adhesion and infiltration during inflammatory processes (Nakajima et al, J. Exp. Med. 1994, 179, 1145-1154). Differential expression of integrins regulates the adhesive properties of cells and different integrins are involved in different inflammatory responses (Butcher et al, Science 1996, 272, 60-66). The α_x integrins, $\alpha_5\beta_1$ (VLA-4) and $\alpha_4\beta_1$ (LPA-1), are expressed primarily on monocytes, lymphocytes, eosinophils, basophils, and macrophages but not on neutrophils (Ellice et al, Cell 1990, 60, 577-584). The primary ligands for α_x integrins are the endothelial surface proteins mucosal addressin cell adhesion molecule (MAdCAM) and vascular cell adhesion molecule (VCAM) with lower affinity (Makarewicz et al, J. Biol. Chem. 1994, 269, 4005-4011). The binding of the $\alpha_5\beta_1$ or $\alpha_4\beta_1$ to MAdCAM and/or VCAM expressed on high endothelial venules (HEVs) at sites of inflammation results in firm adhesion of the leukocyte to the endothelium followed by extravasation into the inflamed tissue (Chuluyan et al, Springer Semin. Immunopathology 1995, 16, 391-404). Monoclonal antibodies directed against $\alpha_5\beta_1$, $\alpha_4\beta_1$, MAdCAM or VCAM have been shown to be effective modulators in animal models of chronic inflammatory diseases such as asthma (Simmons et al, Blood 1992, 80, 388-395), rheumatoid arthritis (RA) (Juliano et al, Current Opinion Cell Biology 1993, 5, 812-818), and inflammatory bowel diseases (IBD) (Lagerge et al, Am. J. Respir. Crit. Care Med. 1995, 151, 822-829 and Barbadillo et al, Springer Semin. Immunopathology 1995, 16). While antibodies have shown efficacy they must be administered parenterally and are inherently cumbersome to produce. Accordingly, it would be desirable to provide small molecule compounds which inhibit the interaction between α_x integrins and ligands MAdCAM and/or VCAM which would be useful for treatment of chronic inflammatory diseases such as arthritis, asthma, multiple sclerosis, Crohn's disease, ulcerative colitis, and hepatitis C.

SUMMARY OF THE INVENTION

According to an aspect of the present invention there is provided compounds of formula (I)

2



wherein

A is a 5 or 6 member, saturated or unsaturated carbocycle or heterocycle optionally substituted by oxo and R_4 .

Q is alkyl, alkenyl or alkynyl optionally substituted with halogen, carboxyl, alkyl or aryl, and wherein one or more carbon atoms are optionally replaced with O, N, NR_6 , S, SO, or SO_2 .

X is $-CR_5-$ or $-N-$.

Y is H, $-CHR_5-$, $-CR_5=$, or a bond.

Z is H, $-CHR_5-$, $-CR_5=$, $-NR_5-$, $=N-$, O, S, SO, SO_2 or a bond, provided that when one of Y and Z is H then the other is also H.

W is $-C(O)NR_6-$, $-NR_6C(O)-$, $-C(S)NR_6-$, $-NR_6C(S)-$, NR_6 , O, S, SO_2 , $-CH_2-$, $-C-$, $-NR_6SO_2-$, $-SO_2NR_6-$, $-OC(O)NR_6-$, $-NR_6C(O)O-$, $-OC(S)NR_6-$, $-NR_6C(S)O-$, $-S-C(S)NR_6-$, $-C(O)-$, $-NR_6C(O)NR_6-$ or $-NR_6C(S)NR_6-$.

R_1 is hydrogen or is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with hydroxyl, halogen, amino, nitro, carboxyl, a carbocycle, or a heterocycle; or R_1 is a carbocycle or heterocycle optionally substituted with hydroxyl, oxo, halogen, amino, or nitro.

R_2 is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with halogen, hydroxyl, oxo, alkoxy, amino, nitro, carboxyl, carboxamide, acyl, acyloxy, amidinyl, guanidinyl, thiol, alkylthio, or one or more carbocycle or heterocycle optionally substituted with halogen, hydroxyl, oxo, alkoxy, amino or carboxyl; or R_2 is a carbocycle or heterocycle optionally substituted with halogen, hydroxyl, oxo, alkoxy, amino, nitro, carboxyl, acyl, acyloxy, alkyl, alkenyl, alkynyl or a carbocycle or heterocycle optionally substituted with halogen, hydroxyl, oxo, alkoxy, amino or carboxyl.

R_3 and R_4 are independently selected from the group consisting of H, hydroxyl, halogen, amino, nitro, carboxyl, alkyl, alkenyl, alkynyl, a carbocycle and a heterocycle, wherein said alkyl, alkenyl, alkynyl, carbocycle and heterocycle groups are optionally substituted with one or more substituents selected from the group consisting of hydroxyl, halogen, amino, oxo and carboxyl, and optionally one or more carbon atoms of said alkyl, alkenyl and alkynyl group is replaced with N, NR_6 , O, S, SO or SO_2 .

R_5 is H or alkyl, alkenyl or alkynyl optionally having a carbon atom replaced with O, N or NR_6 , and optionally substituted with $COOR_7$, or R_5 together with the carbon atom from which it depends forms a double bond to an adjacent carbon or nitrogen atom of Q; or R_5 together with a non-adjacent carbon or nitrogen atom of Q form a carbocycle or heterocycle.

US 6,706,753 B2

3

R_6 is hydrogen, alkyl, alkenyl or alkynyl,
m and n are independently 1, 2 or 3,
and salts, solvates and hydrates thereof


In another aspect of the invention, there is provided pharmaceutical compositions comprising a compound of the invention and a pharmaceutically acceptable carrier, excipient or adjuvant

In another aspect of the invention, there is provided a method of inhibiting binding of an α_v integrin to a protein ligand comprising contacting said α_v integrin with a compound of the invention

In another aspect of the invention, there is provided a method of treating a disease or condition mediated by α_v integrin receptors or ligands of α_v integrin receptors in a mammal, the method comprising administering to said mammal an effective amount of a compound of the invention

DETAILED DESCRIPTION OF THE INVENTION

Compounds are provided having binding affinity for α_v integrins, having the general formula (I)



wherein A, Q, W, X, Y, Z, R_1 to R_6 , m and n are as defined herein

Ring A is a 5 or 6 member, saturated or unsaturated carbocycle or heterocycle optionally substituted by oxo ($=O$) and R_6 . By "carbocycle" is meant herein to be a mono-, bi- or tricyclic ring system containing a 4-16 carbon atom scaffold that is saturated, partially unsaturated or fully unsaturated including aromatic. In the context of ring A, suitable carbocycles include cycloalkyl, cycloalkenyl and aryl. In a preferred embodiment, ring A is a carbocycle selected from the group consisting of cyclopentyl, cyclohexyl and benzene. In a most preferred embodiment ring A is benzene. In another embodiment, ring A is a heterocycle. By "heterocycle" is meant herein to be a mono-, bi- or tricyclic ring system comprising a combination of 4-16 carbon and hetero atoms (i.e. N, O, and S, as well as SO and SO_2) that is saturated, partially unsaturated or fully unsaturated including aromatic. In the context of ring A, preferred heterocycles are 5 and 6 member monocycles. Particularly preferred ring A heterocycles include pyridine, pyran, pyrimidine, pyrazine, pyridazine, pyrrole, furan, thiophene, imidazole, pyrazole, thiazole and triazole. It is appreciated that ring A encompasses heterocycles in which the heteroatoms may be shared with the central ring, if present, and/or may be adjacent to X.

Q is a divalent alkyl, alkenyl or alkynyl linking group optionally substituted with halogen, carboxyl, alkyl or aryl, and wherein one or more carbon atoms are optionally replaced with O, N, NR_6 , S, SO, or SO_2 . In a preferred embodiment, Q is alkyl having 1 to 3 carbon atoms or methylene groups in length, and more preferably length of 2 methylene groups. In another preferred embodiment, the

4

methylene group adjacent to the group X is replaced with a nitrogen atom or NR_6 and more preferably with an oxygen atom. In a particularly preferred embodiment, Q is $-O-CH_2-$ wherein the oxygen atom is adjacent to the X group

X is a bridging group $-CR_5-$ or $-N-$ from which the group Q depends. In a preferred embodiment, X is $-CR_5-$ wherein R_5 is H. In another embodiment R_5 is alkyl, alkenyl or alkynyl optionally having a carbon atom replaced with O, N or NR_6 , and is optionally substituted with $COOR_1$. By "alkyl", "alkenyl" and "alkynyl" is meant herein to be straight or branched aliphatic groups having 1-10 carbon atoms, preferably 1-6 and more preferably 1-4. Preferred alkyl groups are methyl, ethyl, propyl, i-propyl, n-butyl, i-butyl, s-butyl and t-butyl. Preferably, R_5 is an alkyl group having 1 to 3 carbon atoms or methylene groups in length and is substituted with $COOR_1$. More preferably, R_5 is $-(CH_2)_2-COOR_1$. In another embodiment, R_5 together with the carbon atom from which it depends forms a double bond to an adjacent carbon or nitrogen atom of Q. In yet another embodiment R_5 together with a non-adjacent carbon or nitrogen atom from the group Q form a carbocycle or heterocycle. In this context, a preferred embodiment is when R_5 and the carbon atom in Q which is beta to the group X form a 1,3-dioxolane ring. In a particularly preferred embodiment, the dioxolane ring is spiro at X and the two oxygen atoms are alpha to the X group.

Y is H, $-CHR_3-$, $-CR_3-$, or a bond and Z is H, $-CHR_3-$, $-CR_3-$, $-NR_3-$, $-N-$, O, S, SO, SO_2 or a bond, provided that when one of Y and Z is H then the other is also H. In a preferred embodiment, one of Y and Z is a bond while the other is $-CHR_3-$ thereby forming a six member ring fused to ring A and the benzene ring of formula (I) resulting in a tricyclic ring system. In another preferred embodiment Y and Z are both $-CH_2-$ thereby forming a seven member ring. In a more preferred embodiment, Y and Z are both H wherein ring A and the benzene ring of formula (I) are linked via group X rather than forming a fused tricyclic ring system. In a more preferred embodiment, Y and Z are both a bond thereby forming a five member ring fused to ring A and the benzene ring, and in a most preferred embodiment the five member ring together with ring A and the benzene ring of formula (I) form a fluorenyl ring

W is $-C(O)NR_6-$, $-NR_6C(O)-$, $-C(S)NR_6-$, $-NR_6C(S)-$, NR_6 , O, S, SO_2 , $-CH_2-$, $-C-$, $-NR_6SO_2-$, $-SO_2NR_6-$, $-OC(O)NR_6-$, $-NR_6C(O)O-$, $-OC(S)NR_6-$, $-NR_6C(S)O-$, $-S-C(S)NR_6-$, $-C(O)-$, $-NR_6C(O)NR_6-$ or $-NR_6C(S)NR_6-$. In a particular embodiment W is the amido group $-C(O)NR_6-$ wherein the nitrogen atom is adjacent to the benzene ring of formula (I), or alternatively $-NR_6C(O)-$ wherein the carbonyl is adjacent to the benzene ring. In another particularly embodiment W is the sulfonamido group $-NR_6SO_2-$ wherein the sulfur atom is adjacent to the benzene ring, or alternatively $-SO_2NR_6-$ wherein the nitrogen atom is adjacent to the benzene ring. In the context where W is amido or sulfonamido, R_6 is preferably H or C_{1-4} alkyl and more preferably H or methyl. In another particular embodiment, W is O. In a particularly preferred embodiment W is NR_6 wherein R_6 is H or C_{1-4} alkyl and particularly H or methyl

R_1 is hydrogen or is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with hydroxyl, halogen, amino, nitro, carboxyl, a carbocycle, or a heterocycle. By "amino" is meant herein to be a primary, secondary or tertiary amine substituted with alkyl, alkenyl, alkynyl, aryl, or aralkyl each optionally

US 6,706,753 B2

5

substituted as provided herein. By "carboxyl" is meant herein to be $-\text{COOH}$ as well as carboxy ester groups thereof and in particular alkyl esters thereof. Alternatively R_3 is a carbocycle or heterocycle optionally substituted with hydroxyl, oxo, halogen, amino, or nitro. In a preferred embodiment R_3 is H or a group which is liberated in vivo to yield a free carboxy group $-\text{C}(\text{O})\text{O}^-$.

R_2 is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with halogen, hydroxyl, oxo ($=\text{O}$), alkoxy, amino, nitro, carboxyl, carboxamido, acyl, acyloxy, amidinyl ($-\text{C}(\text{NH})\text{NH}-$ or $-\text{NHC}(\text{NH})-$), guanidinyl ($-\text{NHC}(\text{NH})\text{NH}-$), thiol, alkylthio, or one or more carbocycle or heterocycle optionally substituted with halogen (F, Cl, Br or I), hydroxyl, oxo, alkoxy, amino or carboxyl. By "alkoxy" is meant herein to include $-\text{O}-$ alkyl, $-\text{O}-$ alkenyl and $-\text{O}-$ alkynyl wherein alkyl, alkenyl and alkynyl are as previously defined. By "acyl" is meant herein to be a substituted carbonyl ($-\text{C}(\text{O})-$). Suitable acyl groups include alkanoyl, aroyl and aralkanoyl. Suitable acyloxy groups include alkanoyloxy, aroyloxy and aralkanoyloxy. In a particularly preferred embodiment R_2 is a naturally occurring or non-naturally occurring D or L-amino acid residue in which the alpha nitrogen or alpha amino group is optionally acylated. In particularly preferred embodiments, W is NR_6 and R_2 is N-acetyl-tyrosine, D-tyrosine, phenylalanine, benzoyl, isonitrophenyl, 4-methoxyphenylacetyl, 1-fluorenyl-carbonyl, 1-naphthoyl, 2-naphthoyl, 3-hydroxy-phenylalanine, 3-iodo-tyrosine, 3-fluoro-tyrosine, 3-chloro-tyrosine, 4-(4-hydroxyphenyl)-benzoyl, 2,3,5,6-tetrafluoro-tyrosine, 6-hydroxynaphthoyl, 2-phenyl-3-(4-hydroxyphenyl)propanoyl and N-acetyl-3-(4-hydroxyphenyl)proline. In a more preferred embodiment R_2 is a tyrosine amino acid residue and more preferably a tyrosine residue wherein the alpha nitrogen is acylated with acetyl (i.e. N-acetyl tyrosine).

In another embodiment R_2 is a carbocycle or heterocycle optionally substituted with one or more halogen, hydroxyl, oxo, alkoxy, amino, nitro, carboxyl, acyl, acyloxy, alkyl, alkenyl, alkynyl or a carbocycle or heterocycle optionally substituted with halogen, hydroxyl, oxo, alkoxy, amino or carboxyl. In a particular embodiment R_2 is a heterocycle substituted with a phenyl group which in turn is optionally substituted with one or more halogen, hydroxyl, alkoxy or carboxyl. In a particularly preferred embodiment, the heterocycle is a pyrrolidine ring such that when W is an amide group $-\text{C}(\text{O})\text{NR}_6-$ or $-\text{NR}_6\text{C}(\text{O})-$ and W together form a proline amino acid residue. The proline residue is preferably substituted at its beta carbon (i.e. 3-position of the pyrrolidine ring) with a phenyl group which is optionally para-substituted (i.e. at its 4-position) with hydroxyl thereby forming a constrained tyrosine residue. The phenyl group is optionally substituted at one or more of the 2, 3, 5 and 6 positions with a halogen, in particular iodo (I), chloro (Cl) or fluoro (F). Further, the nitrogen atom of the proline residue is optionally acylated, in particular with an alkanoyl group such as acetyl.


R_3 and R_4 are independently selected from the group consisting of H, hydroxyl, halogen, amino, nitro, carboxyl, alkyl, alkenyl, alkynyl, a carbocycle and a heterocycle, wherein said alkyl, alkenyl, alkynyl, carbocycle and heterocycle groups are optionally substituted with one or more substituents selected from the group consisting of hydroxyl, halogen, amino, oxo and carboxyl, and optionally one or more carbon atoms of said alkyl, alkenyl and alkynyl group is replaced with N, NR_6 , O, S, SO or SO_2 . In a particular embodiment R_3 and R_4 are H. Alternatively, one or both of

6

R_3 and R_4 are a carboxy group $-\text{COOR}_1$ linked to the ring from which it depends via a linking group such as an alkyl chain of 1-6 methylene groups in length and preferably 1-3 methylene groups. The linking group may depend from the ring by a functional group such as O (alkoxy), NR_6 (amino), an amido group or a sulfonamido group. In a preferred embodiment, both 'm' and 'n' are the integer 1.

R_6 is hydrogen, alkyl, alkenyl or alkynyl. In a preferred embodiment R_6 is C_{1-4} alkyl and more preferably methyl. In another more preferred embodiment R_6 is H.

In accordance with a preferred embodiment, compounds of the invention have the general formula (II)



wherein

Q is alkyl, alkenyl or alkynyl optionally substituted with halogen, carboxyl, alkyl or aryl, and wherein one or more carbon atoms are optionally replaced with O, N, NR_6 , S, SO, or SO_2 .

R_1 is hydrogen or is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with hydroxyl, halogen, amino, nitro, carboxyl, a carbocycle, or a heterocycle; or R_1 is a carbocycle or heterocycle optionally substituted with hydroxyl, oxo, halogen, amino, or nitro;

R_3 and R_4 are independently selected from the group consisting of H, hydroxyl, halogen, amino, nitro, carboxyl, alkyl, alkenyl, alkynyl, a carbocycle and a heterocycle, wherein said alkyl, alkenyl, alkynyl, carbocycle and heterocycle groups are optionally substituted with one or more substituents selected from the group consisting of hydroxyl, halogen, amino, oxo and carboxyl, and optionally one or more carbon atoms of said alkyl, alkenyl and alkynyl group is replaced with N, NR_6 , O, S, SO or SO_2 .

R_6 is hydrogen, alkyl, alkenyl or alkynyl.

R_7 is hydrogen, hydroxyl, halogen, alkyl, alkoxy or halogen substituted alkyl;

R_8 is H, alkyl, alkenyl or alkynyl.

R_9 is H or $\text{NR}_{11}\text{R}_{12}$, wherein R_{11} and R_{12} are independently H, acyl or an amino acid residue; or one of R_{11} and R_{12} together with R_9 form a heterocycle;

R_{10} is O or S;

m and n are independently 1, 2 or 3,

p is an integer from 1 to 5,

and salts, solvates and hydrates thereof.

In a preferred embodiment R_7 is hydroxyl at the para-position (i.e. 4-position) of the phenyl group. Further R_7 substituents as provided when 'p' is the integer 2, 3, 4 or 5, include 3-iodo, 3-fluoro, 3-chloro and 2,3,5,6-tetrafluoro.

In particular embodiments, R_6 and R_8 are H and R_{10} is oxygen. Alternatively, R_6 is methyl.






In a preferred embodiment, R_9 is $\text{NR}_{11}\text{R}_{12}$, wherein R_{11} is H and R_{12} is H or acyl, in particular alkanoyl i.e. acetyl. In another preferred embodiment, R_9 is $\text{NR}_{11}\text{R}_{12}$, wherein one

US 6,706,753 B2

7







of R_{11} and R_{12} , together with R_6 , form a 5-member hetero-cycle while the other of R_{11} and R_{12} is H or C_{1-4} alkanoyl. Preferably R_{11} and R_6 together form a pyrrolidine ring and R_{12} is H or alkanoyl i.e. acetyl.

Particular compounds of the invention include



8


-continued



US 6,706,753 B2


9

-continued



10


-continued



US 6,706,753 B2


11

-continued



12






-continued



US 6,706,753 B2






13

-continued




14

-continued



US 6,706,753 B2

15



16

lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like.

5 Particularly preferred organic non-toxic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine.

Compounds of the invention may be prepared according to established organic synthesis techniques from starting materials and reagents that are commercially available. In general, the compounds may be prepared starting from a commercially available central ring system. Depending on the particular compound to be prepared, the central ring system is manipulated to append first the carboxylate moiety $-Q-C(O)O-R_1$ to X and then the R_2-W moiety to the benzene ring, or alternatively first appending the R_2-W moiety and then the $-Q-C(O)O-R_1$ moiety.




In a particular embodiment, compounds of formula (Ia), wherein X is C and Q incorporates an oxygen adjacent to X (i.e. an ether), are prepared according to the general scheme 1.

and salts, solvates and hydrates thereof.

It will be appreciated that compounds of the invention may incorporate chiral centers and therefore exist as geometric and stereoisomers. All such isomers are contemplated and are within the scope of the invention whether in pure isomeric form or in mixtures of such isomers as well as racemates. Stereoisomeric compounds may be separated by established techniques in the art such as chromatography, i.e. chiral HPLC, or crystallization methods.

"Pharmaceutically acceptable" salts include both acid and base addition salts. Pharmaceutically acceptable acid addition salt refers to those salts which retain the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, carbonic acid, phosphoric acid and the like, and organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, gluconic acid, lactic acid, pyruvic acid, oxalic acid, malic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, aspartic acid, ascorbic acid, glutamic acid, anthranilic acid, benzoic acid, cinnamic acid, mandelic acid, embonic acid, phenylacetic acid, methane-sulfonic acid, ethane-sulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

Pharmaceutically acceptable base addition salts include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases includes salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, n-propylamine, ethanolamine, 2-diethylaminocethanol, trimethylamine, dicyclohexylamine,




Referring to scheme 1, starting compound (i), wherein A, Y, Z, W, R_2-R_3 , m and n are as previously defined, is reduced with a suitable reducing agent, such as sodium borohydride ($NaBH_4$) or equivalent hydride, to give alcohol (ii). Depending on the particular substituents, starting compound (i) may be commercially available or is otherwise

US 6,706,753 B2

17

prepared according to established organic synthetic techniques from compounds that are commercially available. The final ether compound (1a) of the invention is achieved by treating alcohol (u) with an base such as NaH followed by reacting with electrophilic intermediate (iii), wherein L is a suitable leaving group such as a halogen, i.e. Br, and Q' is the same as Q, i.e. a carboxyl substituted alkyl, alkenyl or alkynyl chain, minus the first carbon atom that would be adjacent to the ring system. It will be appreciated that depending on the particular substituents present in the compound, suitable protection and deprotection procedures will be required as is standard in the art. Numerous protecting groups are described in Greene and Wuts, *Protective Groups in Organic Chemistry*, 2d edition, John Wiley and Sons, 1991, as well as detailed protection and deprotection procedures. For example, suitable amino protecting groups include t-butyloxycarbonyl (Boc), fluorenylmethyloxycarbonyl (Fmoc), 2-trimethylsilyl-ethyloxycarbonyl (Teoc), 1-methyl-1-(4-biphenyl) ethyloxycarbonyl (Bpoc), allyloxycarbonyl (Alloc), and benzyloxycarbonyl (Cbz). Carboxyl groups can be protected as fluorenylmethyl groups and hydroxyl groups may be protected with silyl, monomethoxytrityl, dimethoxytrityl, and trimethoxytrityl groups.

Preparation of compounds of the invention wherein X is carbon and Q is an alkyl chain, may be accomplished according to scheme 2.




Referring to scheme 2, starting compound (i) is reacted with a non-nucleophilic base such as NaH, LDA or TBAF, to give carbanion (ii) which is subsequently reacted with intermediate (iii) to yield compound of formula (1b) of the invention. In a particular embodiment wherein Q is and

18


optionally substituted C_2 alkyl linker to $-\text{COOR}_1$, intermediate carbanion (ii) is reacted with an α, β unsaturated carboxylate (iv) wherein each R is independently hydrogen, halogen, alkyl, carboxyl or aryl, to give compound (1b) of the invention. Preferably R is halogen or C_{1-4} alkyl, i.e. methyl.

In another embodiment, compounds of formula (I), wherein X is N, are prepared according to scheme 3



Referring to scheme 3, starting compound (i) is reacted with intermediate (ii) wherein L is a suitable leaving group, i.e. Br, in the presence of a non-nucleophilic base such as Ca_2CO_3 or K_2CO_3 to yield the final compound (1c) of the invention. Alternatively, starting compound (i) can be reacted with an α, β unsaturated carboxylate (iii) as in scheme 2 to give compound (1c) of the invention.


In another embodiment, compounds of formula (I) wherein W is $-\text{C}(\text{O})\text{NH}-$, $-\text{SO}_2\text{NH}-$ or $-\text{NH}-$ are prepared according to scheme 4.



US 6,706,753 B2

19

-continued




Referring to scheme 4, from the starting amine compound (i) may be prepared amide (Id), sulfonamide (Ie), amine (If), carbamate (Ig) and urea (Ih) by reacting (i) respectively with intermediates (ii) an activated ester of R_2-CO_2H , (iii) R_2-SO_2-L wherein L is a leaving group such as a halogen, (iv) R_2-L in the presence of a non-nucleophilic base or $R_2-C(O)H$ in the presence of $NaCNBH_3$, (v) $R_2-OC(O)-L$, and (vi) isocyanate $R_2-N=C=O$. It will be appreciated that thioamides may be prepared in a similar manner as the

20


amides by using $R_2-C(S)H$ or $R_2-C(S)O-$, and thiocarbamates may be prepared using $R_2-S-C(O)-L$ or $R_2-OC(S)-L$; and thioureas may be prepared using isothiocyanate $R_2-N=C=S$. In a particular

5 In another embodiment, compounds of formula (I) wherein W is $-O-$ or $-S-$ may be prepared according to scheme 5a and 5b.


scheme 5a



scheme 5b




US 6,706,753 B2





Referring to schemes 5a and 5b, the starting hydroxyl or thiol compound is reacted with a non-nucleophilic base such as NaH to remove a proton. The anion is then reacted with R_1-L wherein L is a suitable leaving group to yield the resulting ether (II) and thioether (Ij) compound of the invention.

In another embodiment compounds of formula (I), wherein W is carbonyl $-C(O)-$, may be prepared according to scheme 6.




Referring to scheme 6, starting compound (i) is reacted with aldehyde intermediate (ii) in the presence of NaCNBH₃ to give compound (Ik) of the invention.

In another embodiment, compounds of formula (I), wherein W is $-NR_2SO_2-$, are prepared according to scheme 7.



Referring to scheme 7, starting compound (i) is reacted with sulfonic acid and then POCl₃ to give the sulfonyl chloride compound (iii) which is then reacted with R_2-NH_2 to give compound (II) of the invention.


In another embodiment, compounds of formula (I), wherein W is C, may be prepared according to scheme 8.



US 6,706,753 B2


23

-continued



Referring to scheme 8, starting compound (i) is reacted with Pd^0 catalyst followed by $\text{R}_2\text{---B(OH)}_2$ or $\text{R}_2\text{---C}\equiv\text{CH}$ to give final compound (Im) or (In) respectively. In another embodiment, compounds of formula (I), wherein W is carbamate $\text{---NR}_4\text{C(O)O---}$ or thiocarbamate $\text{---NR}_4\text{C(S)O---}$, may be prepared according to scheme 9.

scheme 9




Referring to scheme 9, starting alcohol compound is reacted with intermediate $\text{R}_2\text{---NC(O)-L}$ or $\text{R}_2\text{---NC(S)-L}$, wherein L is a suitable leaving group such as Br, to give final compound of the invention (Io) or (Ip) respectively. In a particular embodiment, a morpholino carbamate compound of the invention by reacting the starting alcohol with morpholine- C(O)-Cl .

In another embodiment, compounds of formula (I), wherein W is thiocarbamate $\text{---NR}_4\text{C(O)S---}$ or thiocarbamate $\text{---NR}_4\text{C(S)S---}$, may be prepared according to scheme

24

scheme 10



Referring to scheme 10, starting thiol compound (i) is reacted with intermediate $\text{R}_2\text{---NC(O)-L}$ or $\text{R}_2\text{---NC(S)-L}$, wherein L is a suitable leaving group such as Br, to give final compound of the invention (Iq) or (Ir) respectively.

In an aspect of the invention, there is provided a method of inhibiting binding of an α_v integrin to a ligand, the method comprising contacting said α_v integrin with a compound of formula (I). The method may be carried out as a solution based or cell based assay wherein the compound of the invention is introduced to the integrin in the presence of a putative or known ligand of the integrin. The compound may be labeled, for example isotopically radiolabeled, to facilitate detection of ligand binding or lack thereof to the integrin. Thus compounds of the invention are useful for diagnostic assays.

Compounds of the invention are useful to prevent the interaction of an epithelial cell bearing VCAM-1 and/or MAdCAM on the cell surface with a leukocyte cell bearing $\alpha_4\beta_1$ and/or $\alpha_4\beta_7$ on the surface by contacting the epithelial cell or the leukocyte with an inhibitory amount of the compound of the invention. The compounds are useful in assays to determine the inhibitory effect of a compound which antagonizes the binding of $\alpha_4\beta_1$ and/or $\alpha_4\beta_7$ integrin to VCAM-1 ligand and/or MAdCAM ligand. The inhibitory compound may be a small molecule, a protein or peptide or an antibody. In an in vitro assay, the ligand or the integrin may be directly or indirectly bound to a surface, such as microtiter plate, using known methods described for example in WO 9820110, WO 9413312, WO 9624673, WO 9806248, WO 9936393, and WO 9910312. The other member of the binding pair, e.g. the integrin or the ligand, respectively, (or a cell expressing the same on its surface) is

US 6,706,753 B2

25

then added to the surface bound member and the inhibitory effect of a test molecule is determined. The inhibitory activity of the compounds of the invention can also be determined with this type of assay.

The binding of the integrins to their respective ligands is known to be involved in inflammatory conditions associated with leukocyte infiltration of tissues lined with epithelial cells expressing VCAM-1 or MADCAM. Such tissues include the gastrointestinal tract, skin, urinary tract, respiratory airways and joint synovial tissues. The compounds of the invention are useful in treating diseases in which such binding is implicated as a cause of the disease or symptoms of the disease. Undesired disease symptoms may arise from cell adhesion and/or cell activation which releases proinflammatory mediators, typically when there is an increase or upregulation in the expression of VCAM-1 and/or MADCAM on the surface of endothelial cells. Various disease states which can be treated and for which the inflammatory symptoms can be reduced upon administration of the compounds of the invention include rheumatoid arthritis, asthma, psoriasis, multiple sclerosis, inflammatory bowel disease including ulcerative colitis, pouchitis and Crohn's disease, Celiac disease, non-tropical Sprue, graft-versus-host disease, pancreatitis, insulin-dependent diabetes mellitus, mastitis, cholecystitis, pericholangitis, chronic sinusitis, chronic bronchitis, pneumonitis, collagen disease, eczema, and systemic lupus erythematosus. Compounds of the invention are useful in treating these diseases and conditions by inhibiting the integrin/ligand binding. Compounds of the invention are therapeutically and/or prophylactically useful for treating diseases or conditions mediated by α , integrin receptors i.e. $\alpha_v\beta_3$ and $\alpha_v\beta_1$, and/or their ligands, in particular VCAM-1 and MADCAM-1. Accordingly in an aspect of the invention, there is provided a method of treating a disease or condition mediated by the α_v integrin receptors or ligands of α_v integrin receptor ligands in a mammal, for example a human, comprising administering to said mammal an effective amount of a compound of the invention. By "effective amount" is meant an amount of compound which upon administration is capable of reducing the amount of ligand able to bind to α_v integrins in vivo, or an amount of compound which upon administration is capable of alleviating or reducing the severity of symptoms associated with the disease or condition mediated by α_v integrins or ligands thereof.

The actual amount of compound administered and the route of administration will depend upon the particular disease or condition as well as other factors such as the size, age, sex and ethnic origin of the individual being treated and is determined by routine analysis. In methods of the invention, the compound may be administered orally (including buccal, sublingual, inhalation), nasally, rectally, vaginally, intravenously, intradermally, subcutaneously and topically. Compounds will be formulated into compositions suitable for administration for example with suitable carriers, diluents, thickeners, adjuvants etc as are routine in the formulation art. Accordingly, another aspect of the invention provides pharmaceutical compositions comprising a compound of formula (I) and a pharmaceutically acceptable carrier, excipient or adjuvant. Compositions of the invention may also include additional active ingredients. Dosage forms include solutions, powders, tablets, capsules, gel capsules, suppositories, topical ointments and creams and aerosols for inhalation. Formulations for non-parenteral administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable adjuvants. Pharmaceutically acceptable organic or inorganic car-

26

rier substances suitable for non-parenteral administration which do not deleteriously react with compounds of the invention can be used. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with compounds of the invention. Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. Optionally, the suspension may also contain stabilizers.

In a preferred embodiment, compounds of the invention are administered via oral delivery. Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, troches, tablets or SECs (soft elastic capsules or caplets). Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids, carrier substances or binders may be desirably added to such formulations. Such formulations may be used to affect delivering the compounds to the alimentary canal for exposure to the mucosa thereof. Accordingly, the formulation can consist of material effective in protecting the compound from pH extremes of the stomach, or in releasing the compound over time, to optimize the delivery thereof to a particular mucosal site. Enteric coatings for acid-resistant tablets, capsules and caplets are known in the art and typically include acetate phthalate, propylene glycol and sorbitan monolactate.

Various methods for producing formulations for alimentary delivery are well known in the art. See, generally *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, Pa., 1990. The formulations of the invention can be converted in a known manner into the customary formulations, such as tablets, coated tablets, pills, granules, aerosols, syrups, emulsions, suspensions and solutions, using inert, non-toxic, pharmaceutically suitable excipients or solvents. The therapeutically active compound should in each case be present in a concentration of about 0.5% to about 99% by weight of the total mixture, that is to say in amounts which are sufficient to achieve the desired dosage range. The formulations are prepared, for example, by extending the active compounds with solvents and/or excipients, if appropriate using emulsifying agents and/or dispersing agents, and, for example, in the case where water is used as the diluent, organic solvents can be used as auxiliary solvents if appropriate.

Compositions may also be formulated with binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrates (e.g., starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulfate). Tablets may be coated by methods well known in the art. The preparations may also contain flavoring, coloring and/or sweetening agents as appropriate.


Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, sachets or tablets each containing predetermined amounts of the active ingredients; as powders or granules; as solutions or suspensions in an aqueous liquid or a non-

US 6,706,753 B2

27

aqueous liquid, or as oil-in-water emulsions or water-in-oil liquid emulsions. A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine, the active ingredients in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredients therein.

EXAMPLE 1
Synthesis of Compound 5



28

Commercially available 2-amino-9-fluorenone 1 (10 mmol) was dissolved in 100 mL of methanol and 0.38 g (10 mmol) of sodium borohydride was added. The mixture was stirred at 25° C for 1 h and concentrated to dryness under vacuum. The residue was suspended in 100 mL of saturated aqueous NaHCO_3 and extracted with ethylacetate (100 mL). The ethylacetate layer was concentrated under vacuum and the residue was crystallized from ethylacetate/hexane (2:1) to afford 1.4 g of pure 2-amino-9-hydroxyfluorene 2.


Compound 2 (0.39 g, 2 mmol) was dissolved in 50 mL of anhydrous DMF and cooled to 0° C. under nitrogen. Lithium bis(trimethylsilyl)amide (2 mmol; 1M in THF) was added dropwise and stirring was continued for 30 min. Iodoethylacetate (3 mmol) was added dropwise and the solution was stirred 12 h at 0° C. The reaction mixture was poured into 200 mL of water and extracted with ethyl acetate (200 mL). The extract was concentrated and the product was purified via silica gel chromatography (25% ethylacetate/hexane) to afford 0.34 g of fluorenyl ether 3.

Fluorenyl ether 3 (0.28 g; 1 mmol) was dissolved in 5 mL of DMF and N-Fmoc-L-tyrosine (1 mmol) was added followed by HATU (1 mmol) and N-methylmorpholine (2 mmol). The reaction was stirred overnight at 25° C., poured into 50 mL water, and the product extracted with ethylacetate. Concentration under vacuum afforded 0.5 g of crude 4 which was used directly in the next step without further purification.

Compound 4 (0.5 g) was dissolved in 1:1 THF/water and 3 mL LiOH (1M in water) was added. The reaction was stirred for 4 h at 25° C. and concentrated under vacuum to remove most of the THF. The aqueous solution was lyophilized and the crude product was purified by reverse HPLC (acetonitrile/water gradient) to afford 90 mg of pure 5 as the TFA salt.

EXAMPLE 2


Synthesis of 2-Amino Carbazole



US 6,706,753 B2

29

-continued




Carbazole (100.24 g) was suspended in acetic anhydride (300 mL) with catalytic boron trifluoride etherate (0.65 mL) and the solution refluxed for 25 min, then cooled to 0° C. and the solid was then collected and recrystallized from hexane to give 82.65 g 9-acetyl-carbazole.

9-Acetyl-carbazole (41.85 g, 0.2 mol) was dissolved in 1 L methylene chloride and 28.5 mL acetylchloride (0.4 mol) and 120 g aluminum chloride (0.9 mol) were added and the solution refluxed for 1.5 h. The solution was cooled to -78° C. and 6N HCl was added slowly while stirring and allowed to warm to room temperature. Methylene chloride was added to dissolve the precipitate and the solution was extracted, and the organics dried over Na₂SO₄, decolorized with charcoal, filtered and concentrated. Recrystallization from benzene/hexane gave 27.9 g 2,9-diacetylcarbazole.


2,9-Diacetylcarbazole, (25.1 g, 0.1 mol) was dissolved in 100 mL pyridine. Hydroxylamine hydrochloride (10.42 g, 0.15 mol) was added and the solution refluxed for 20 min. The mixture was cooled and poured into 70 mL conc HCl in ice. The precipitate was collected by filtration, washed with a large volume of water and dried in vacuum over KOH to yield 24 g white solid. 5 g of this solid (18.8 mmol) was dissolved in 17 mL TFA and 0.3 mL trifluoroacetic anhydride (2.1 mmol) was added. The mixture was refluxed for 20 min then concentrated to give the crude 2,9-diacetylcarbazole amine. Both acetyl groups were removed by suspending the crude material in 400 mL 10% aq. KOH and refluxing for 1 day. The solution was cooled and ethyl acetate added and the product 2-aminocarbazole was extracted, washed with brine, dried over Na₂SO₄ and concentrated.

EXAMPLE 3

Synthesis of Compound 6



1) Ac-Tyr
HBTU/DMF
2) Ethyl bromoacetate
C₂CO₃
3) LiOH




To 0.1 g (0.55 mmol) 2-aminocarbazole in 6 mL DMF was added 0.37 g (1.65 mmol) of N-Ac-L-Tyr, 0.37 g (1.65

30


mmol) HBTU and 0.3 mL (1.7 mmol) DIPEA. The reaction was stirred for 1 h, poured into ethyl acetate and washed consecutively with 0.1N H₂SO₄, aq. NaHCO₃ and brine then dried over Na₂SO₄ and concentrated. Flash chromatography (10% methanol/methylene chloride) gave 0.16 g (0.26 mmol) of the acetylated amine. This was stirred for 1 h with cesium carbonate (0.185 g, 0.57 mmol) and ethyl bromoacetate (0.043 mL, 0.39 mmol) in 4 mL DMF and then poured into ethyl acetate and washed with saturated NH₄Cl and brine and dried over Na₂SO₄ and concentrated. Without purification this was dissolved in 5 mL THF and 5 mL water and 0.043 g (1 mmol) LiOH was added. The ester was cleaved in <30 min and then diluted with ethyl acetate and washed consecutively with 0.1N H₂SO₄ and brine and then dried over Na₂SO₄ and concentrated. The product was lyophilized and purified by HPLC.

EXAMPLE 4

Synthesis of Compound 7



1) const. N-Ac-Tyr
HBTU/DIPEA
2) Ethyl bromoacetate
C₂CO₃
3) LiOH



To 6.37 g (31.44 mmol) of the 9-acetylcarbazole in 200 mL CH₂Cl₂ was added 10.68 g silver nitrate (63 mmol) and 25.1 g aluminum chloride (188.6 mmol). This was stirred at room temperature for 2 h, diluted with methylene chloride and poured onto ice containing 82 mL conc HCl. The organic layer was separated, washed three times with water, dried with MgSO₄ and concentrated to a dark brown solid. The solid was dissolved in 200 mL methanol, 4% aq. KOH was added and the solution was refluxed for 1 h. The solution was cooled and concentrated, then taken up in ethyl acetate and washed consecutively with water and brine then dried over Na₂SO₄. The 9-nitrocarbazole (0.1 g, 0.47 mmol) was dissolved in 3 mL DMF and 0.13 mL (1.42 mmol) methyl acrylate and 0.69 g (2.12 mmol) C₂CO₃ were added. The reaction was stirred for 2 h, diluted with ethyl acetate and washed with saturated NH₄Cl and brine and then dried over Na₂SO₄ and concentrated. Flash chromatography (25% ethyl acetate/hexane) gave 0.081 g alkylated 9-nitrocarbazole. This was dissolved in 5 mL methanol and a spatula of Pd/C and a balloon of H₂ were used to reduce the nitrocarbazole to the amino carbazole. This was dissolved in 5 mL DMF and 0.167 g (0.75 mmol) N-Ac-L-Tyr, 0.394 g (0.96 mmol) HBTU and 0.19 mL (1.12 mmol) DIPEA were added. This was stirred for 1 h, diluted with ethyl acetate and washed consecutively with 0.1N H₂SO₄, aq. NaHCO₃ and brine and then dried over Na₂SO₄ and


US 6,706,753 B2

31

concentrated. Without purification this was dissolved in 5 mL THF and 5 mL water and 0.063 g (1.49 mmol) LiOH was added. The ester was cleaved in <20 min and diluted with ethyl acetate and washed consecutively with 0.1N H₂SO₄ and brine. This was dried over Na₂SO₄ and concentrated and then lyophilized and purified by HPLC.

EXAMPLE 5


Synthesis of Compound 8




To 0.3 g (1.64 mmol) 2-aminocarbazole in 6 mL DMF was added 1.2 g (6.59 mmol) of the constrained-N-Ac-L-Tyr, 2.31 g (6.1 mmol) HBTU and 2 mL (11.5 mmol) DIPEA. The reaction was stirred for 2.5 h, poured into ethyl acetate and washed consecutively with 0.1N H₂SO₄, aq. NaHCO₃ and brine and then dried over Na₂SO₄ and concentrated. Flash chromatography (10% methanol/methylene chloride) gave 0.60 g of the acrylated amine. 0.20 g of this was stirred for 1 h with cesium carbonate (0.349 g, 1.07 mmol) and ethyl bromoacetate (0.08 mL, 0.73 mmol) in 5 mL DMF which was then poured into ethyl acetate and washed with saturated NH₄Cl and brine and then dried over Na₂SO₄ and concentrated. Without purification this was dissolved in 4 mL THF and 4 mL water and 0.082 g (1.95 mmol) LiOH was added. The ester was cleaved in <30 min and then diluted with ethyl acetate and washed consecutively with 0.1N H₂SO₄ and brine. This was then dried over Na₂SO₄, concentrated and then lyophilized and purified by HPLC.

EXAMPLE 6

Synthesis of Compound 9



32



10

15


20

25

30

EXAMPLE 7

Synthesis of 3-Amino-Dibenzosuberol



35

40

45

50

55

60

65

Dibenzosuberol (6.0 g, aldrich) was added to 100 ml of stirred concentrated nitric acid at ambient temperature. Periodic reaction progress analysis (i.e., NMR and IR) indicated that dibenzosuberol was first oxidized to dibenzosuberone, then nitrated at the 3-position. After 2 hours, the reaction mixture was poured into ice water and extracted with 1:1 hexane:ethyl acetate. The crude product was purified by

US 6,706,753 B2


33

Flash chromatography on silica (9:1 hex/EtOAc) to yield 3.22 g (45%) of 3-amino-5-(ethyl carboxymethyl) dibenzosuberone.

3-mirobenzosuberone (3.22 g) was reduced with hydrogen at 40 psi in 1:1 methanol/THF (40 ml), acetic acid (2 ml) and 10% Pd/C (0.5 g) for 1.5 hours. Filtration and concentration gave 3-aminodibenzosuberone 1.35 g (48%). 3-aminodibenzosuberone (1.35 g) and sodium borohydride (3 g) were dissolved in 100 ml THF. Ethanol (30 ml) was added in portions over one hour with stirring at ambient temperature. After 18 hours, 1N HCl was added dropwise until a pH of 7 was obtained. The reaction mixture was partitioned between 1:1 hexane/ethyl acetate and saturated aqueous sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to give 1.0 g of crystalline 3-aminodibenzosuberol (74%).

EXAMPLE 8

Synthesis of Compound 10



3-aminodibenzosuberol (0.95 g) was dissolved in dry DMF (10 ml). Sodium hydride (0.25 g of 60% in oil) was added and the reaction mixture stirred until hydrogen evolution ceased. Ethyl iodacetate (1.35 g) was added and the reaction mixture stirred at ambient temperature for 3 hours then partitioned between ethyl acetate and water, washed with brine, dried and concentrated. Flash chromatography on silica (75:25, hex:EtOAc) gave 0.79 g of 3-amino-5-(ethyl carboxymethyl) dibenzosuberone (60%).


3-amino-5-(ethyl carboxymethyl) dibenzosuberone (395 mg), N-acetyl tyrosine (425 mg), HBTU (723 mg), diisopropylethylamine (491 mg) and 5 ml anhydrous DMF were combined and stirred at ambient temperature overnight. The reaction was partitioned between ethyl acetate and 10% aqueous citric acid. The organic phase was dried, filtered and the product purified on silica (96:4 methylene chloride:methanol) to give 290 mg of 24015-94.

The above product (24015-94, 50 mg) was dissolved in 8 ml of 1:1 ethanol/water and 6.1 mg of lithium hydroxide monohydrate added. After 3 hours, hydrolysis was complete and 1 ml of acetic acid was added. The reaction mixture concentrated and the product purified by HPLC (1 inch C-18, 10 to 90 in 60 min, 254 nm).

34

EXAMPLE 9

Synthesis of Compound 11



68 μ L of acrylic acid (0.25 mmol, R=H) was added to 4 mL of DCM followed by 135 mg of HOBt (0.25 mmol) and 157 μ L of DIPEA (0.25 mmol). After equilibration for 2 min

US 6,706,753 B2

35

this solution was combined with 200 mg of Wang polystyrene resin (0.5 mmol/g, 0.1 mmol) and allowed to shake for 4 h at which time the resin was filtered and washed extensively with repeating volumes of DMF, THF, MeOH, and DCM. In a dry vial, 2 mL of THF was combined with 2 mL of a 1.0 M solution of TBAF in THF (Aldrich Chemical Company). 362.5 mg of 2-aminofluorene (X=H) was added to the TBAF solution and allowed to shake for 10 min prior to adding it to the previously mentioned resin. The resin was stirred for 16 h and the washed as before. At this point, the amino acid tyrosine was coupled to the amine on the fluorenyl ring by treating the resin with 4 mL of a 0.25 M solution of the amino acid, HOAt, HATU, and DIPEA in DMF for 16 h. After washing the resin the FMOC protecting group was removed by mixing the resin with 4 mL of a 25% piperidine solution in NMP for 10–15 min. The piperidine was removed from the resin with repeated washings and the acylation was performed on the resin using a solution of acetyl anhydride and DIPEA each at a 0.25 M concentration in 4 mL of DCM for 1 h, followed by a washing step. The compound was liberated from the resin by treating it with 4 mL of a 90% TFA, 5% DCM, 2.5% water and 2.5% triethylsilane cocktail for 1 h. The TFA solution was filtered from the resin and evaporated under a stream of nitrogen gas. The product was purified on a C-18 reversed phase HPLC to yield 3 mg of product (R, R'-H) and confirmed by mass spectroscopy.

The reaction with the acrylate could also be performed where R was an alkyl group. In cases where R'-Br then Suzuki couplings using aryl boronic acids were performed prior to the acylation with FMOC-tyrosine. A typical procedure is as follows. The resin (100 mg) is swelled in 2 mL NMP containing 0.25 M boronic acid, 0.25 M DIPEA, and 50 mg of Palladium catalyst, Pd(PPh₃)₄Cl₂, and heated to 60° C. for 6 hours. A variety of amine or carbonate bases and palladium catalysts could be also used to perform this transformation. Additionally the nitrogen on the tyrosine ring could be acylated or sulfonylated under a variety of conditions to yield products with functional groups other than the acetyl group. By products resulting from an additional acrylate adding to the 9-position of the fluorenyl ring were observed with higher loading resins and in some cases were isolated in significant quantities.

EXAMPLE 10

Biological Activity Assays

The compounds of the invention can be assayed for ability to block the $\alpha_v\beta_3$ /MAdCAM-1 or $\alpha_v\beta_1$ /VCAM-1 binding interaction by addition of serial dilutions of the samples to plates with the receptors as follows. 96-well plates are coated with mouse anti-human α_v (31470D, Pharmingen, San Diego, Calif.). The plates are decanted and blocked with 0.5% BSA. After washing $\alpha_v\beta_1$ or $\alpha_v\beta_3$ is added, followed by incubation for 2 h at room temperature. The plates are washed and samples of the small molecule antagonists are added to the plates with MAdCAM-1-Ig-HRP or VCAM-1-Ig-HRP for 2 h at room temperature. After an additional wash, the bound MAdCAM-1-Ig-HRP or VCAM-1-Ig-HRP is detected by addition of tetramethylbenzidine (TMB, Kirkegaard & Perry, Gaithersburg, Md.), followed by detection of the absorbance of the product.

Alternatively, the compounds can be assayed using any known protein-protein or cell-based assay method, such as those described, for example, in WO 99/10312 (examples 179–180) and WO 99/36393 (RPMI-CS-1 cell adhesion assay). See also Cardarelli et al., 1994, J. Biol. Chem., 269:18668–18673.


36

For example, 96-well ELISA plates are coated overnight at 4° C with 2 μ g/ml with anti-human CD49d, (31470D, Pharmingen, San Diego, Calif.) in phosphate buffered saline. The plates are decanted and blocked with assay buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM MnCl₂, 0.05% Tween-20 and 0.5% BSA) at room temperature for one hour, with gentle shaking. The plates are washed three times (in 50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 1 mM MnCl₂, 0.05% Tween-20) and 2 μ g/ml of the desired integrin in assay buffer is added, followed by incubation at room temperature for two hours, with gentle shaking. After washing three times, 50 μ l of samples of the small molecule antagonists (serial dilutions from 10 mM stocks in 100% DMSO) are added to the plates with 50 μ l of 1 μ g/ml MAdCAM-1-Ig-HRP or VCAM-1-Ig-HRP in assay buffer. The plates are incubated two hours at room temperature, with gentle shaking, followed by washing six times. The bound MAdCAM-1-Ig-HRP or VCAM-1-Ig-HRP is detected by addition of the peroxidase substrate, 3, 3', 5, 5'-tetramethylbenzidine (TMB, Kirkegaard & Perry, Gaithersburg, Md.), for 10 minutes, followed by addition of 1M phosphoric acid to stop the reaction. The absorbance of the solutions are read at 450 nm on a plate reader.

Suitable animal models exist for many diseases and conditions which can be treated with the compounds of the invention. Additional confirmation of the efficacy of these compounds in specific diseases and at desired doses can be assayed using these established models. For example, animal models of chronic inflammatory diseases such as asthma (Laberge, S. et al., Am. J. Respir. Crit. Care Med., 1995, 151:822–829), rheumatoid arthritis (RA, Barbadillo, C. et al., Springer Semin. Immunopathol., 1995, 16:375–379), and inflammatory bowel diseases (IBD; Podalski, D. K., N. Eng. J. Med., 1991, 325:928–937; Powrie, F. et al., Ther. Immunol., 1995, 2:115–123) may be used to demonstrate the activity of the compounds of the invention and to conduct dose and efficacy studies.

We claim:

1. A compound having the general formula (II)



wherein

is alkyl, alkenyl or alkynyl optionally substituted with halogen, carboxyl, alkyl or aryl, and wherein one or more carbon atoms are optionally replaced with O, N, NR₂, S, SO, or SO₂;

R₁ is hydrogen or is selected from the group consisting of alkyl, alkenyl and alkynyl, each of which is optionally substituted with hydroxyl, halogen, amino, nitro, carboxyl, a carbocycle, or a heterocycle; or R₁ is a carbocycle or heterocycle optionally substituted with hydroxyl, oxo, halogen, amino, or nitro,

R₃ and R₄ are independently selected from the group consisting of H, hydroxyl, halogen, amino, nitro,

US 6,706,753 B2

37

carboxyl, alkyl, alkenyl, alkynyl, a carbocycle and a heterocycle, wherein said alkyl, alkenyl, alkynyl, carbocycle and heterocycle groups are optionally substituted with one or more substituents selected from the group consisting of hydroxy, halogen, amino, oxo and carboxyl, and optionally one or more carbon atoms of said alkyl, alkenyl and alkynyl group is replaced with N, NR₆, S, SO or SO₂;

R₆ is hydrogen, alkyl, alkenyl or alkynyl;

R₇ is hydrogen, hydroxyl, halogen, alkyl, alkoxy or halogen substituted alkyl,

R₈ is H, alkyl, alkenyl or alkynyl;

R₉ is NR₁₁R₁₂, wherein one of R₁₁ and R₁₂ is H, acyl or an amino acid residue, and the other together with R₉ forms a heterocycle;

R₁₀ is O or S,

m and n are independently 1, 2 or 3,

p is an integer from 1 to 5,

and salts, solvates and hydrates thereof.

2 The compound according to claim 1, wherein Q is an alkyl chain of 1 to 3 carbon atoms in length.

3 The compound according to claim 2, wherein said alkyl chain is 2 carbon atoms in length.

4 The compound according to claim 2, wherein one of the carbon atoms in said alkyl chain is replaced with an oxygen atom.

5 The compound according to claim 4, wherein said oxygen atom is adjacent to the ring.


6 The compound according to claim 1, wherein R₁ is H.

7 The compound according to claim 1, wherein R₃ and R₄ are both H.





8 The compound according to claim 1, wherein R₆ is H.

9 The compound according to claim 1, wherein p is the integer 1, and R₇ is OH at the para-position.

10 A compound according to claim 1 selected from the group consisting of



38



and salts, solvates and hydrates thereof